

Award Number: W81XWH-11-2-0165

TITLE: Linking NASA Data with Environmental Exposures and Health Outcomes in Theater of War

PRINCIPAL INVESTIGATOR: David Lary

CONTRACTING ORGANIZATION: University of Texas at Dallas  
Richardson, TX 75080

REPORT DATE: October 2013

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>					
1. REPORT DATE October 2013		2. REPORT TYPE Final		3. DATES COVERED 23 September 2011- 22 September 2013	
4. TITLE AND SUBTITLE Linking NASA Data with Environmental Exposures and Health Outcomes in Theater of War				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-11-2-0165	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) David Lary  david.lary@utdallas.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Texas at Dallas Richardson, TX 75080				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Numerous studies have associated acute and chronic exposures to high levels of particulate matter(PM <sub>10/2.5</sub> ) with health outcomes such as increased hospital admissions, increased respiratory/circulatory symptoms, and decreased lung functions. These exposures, which come from a variety of sources such as blowing sand and dust, smoke, vapors, and aerosols, are common in many areas throughout the globe where U.S. military personnel are deployed in support of our national defense.					
15. SUBJECT TERMS- none provided					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT  UU	18. NUMBER OF PAGES  16	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

## Table of Contents

	<u>Page</u>
Introduction.....	4
Hypothesis.....	4
Technical Objectives.....	5
Method.....	6
References.....	12

## Introduction

Numerous studies have associated acute and chronic exposures to high levels of particulate matter ( $PM_{10/2.5}$ ) with health outcomes such as increased hospital admissions, increased respiratory/circulatory symptoms, and decreased lung functions. These exposures, which come from a variety of sources such as blowing sand and dust, smoke, vapors, and aerosols, are common in many areas throughout the globe where U.S. military personnel are deployed in support of our national defense.

Addressing such health concerns requires an accurate assessment of the small-aerosol-particle concentration near the ground—for example, the issue of air pollution and its effects on health. Various networks of ground-based sensors provide routine measurements of  $PM_{2.5}$ , but their spatial coverage is rather sparse, especially in Third World countries. On the other hand, multiple space-borne sensors measure the total aerosol optical depth (AOD), often with nearly global daily coverage. However, while we have good global coverage for AOD, it is often not the best proxy for the near-surface aerosol concentration. This can be for a variety of reasons, for example, aerosols can be transported at high altitudes without being present near the land surface, while still contributing to a high total AOD. While there is a growing interest in using satellite data, there is the issue that the currently available satellite data products do not provide accurate data on the near-surface  $PM_{2.5}$  abundance that we need for health studies (Hoff and Christopher 2009).

We propose to break through these limitations by bringing together data from multiple sensors and new machine learning methodology. We will also be using new methodology that has recently won recognition as a NASA Aura mission science highlight and the 2010 IEEE Geoscience and Remote Sensing Society Letters Prize Paper Award. This methodology takes into account the cardinally nonlinear relationship between the near-surface abundance of  $PM_{2.5}$  and AOD, which is a function of the boundary-layer height, humidity, surface pressure, surface wind speed, and surface type. This is a major achievement, as, although we have numerous observations, we do not yet have a complete theoretical understanding of this cardinally nonlinear relationship between the near surface abundance of  $PM_{2.5}$  and AOD.

## Hypothesis

The hypothesis of this proposal is that a suite of remote sensing data products on atmospheric aerosols used in their meteorological context and processed by machine learning can provide a daily estimate of the global  $PM_{2.5}$  abundance. This information is of considerable value to Global Health Surveillance (GHS), providing a capability to routinely estimate troop deployment exposure to elevated levels of

particulate matter (PM) globally, significantly contributing to DoD-wide force health protection initiatives.

## Technical Objectives

*The goal of this study is to provide a quantitative understanding of the intrinsically nonlinear, multivariate relationship between the abundance of PM<sub>2.5</sub> in the atmospheric boundary layer and Remotely Sensed Aerosol Optical Depth (AOD) and extinction products. This is encapsulated in a software system that is capable of routinely providing a global data product for DoD health applications. As this project nears completion we see that we the basis for an operational system to serve the DoD health system. This can provide global coverage and therefore has the potential to provide Global Health Surveillance (GHS) with a capability to routinely estimate troop deployment exposure to elevated levels of PM globally, significantly contributing to DoD-wide force health protection initiatives.*

Realizing our goal required two components. The first is to use the appropriate temporally and spatially varying meteorological *context* of the latest version of each satellite product, as well as *in-situ* ground truth observations of PM<sub>2.5</sub> abundance. The precise context of observations is critically important, as there is significant temporal and spatial variability in the abundance of PM<sub>2.5</sub>, so careful attention must be paid to ingesting/fusing the satellite observations at both the appropriate time and place.

The second required component uses nonlinear, nonparametric, multivariate machine learning to address the issues for which we do not yet have a complete theoretical description encapsulated in our Numerical Weather Prediction (NWP) models. It would obviously be ideal if we had a complete theoretical understanding of the multivariate, nonlinear relationship between PM<sub>2.5</sub> and AOD, in which case we would gladly dispense with the machine learning. However, as this most desirable state currently eludes us, the array of tools we have for multivariate, nonlinear, nonparametric machine learning has proved invaluable to a wide variety of applications and has already won significant recognition within NASA. In this study the nonlinear, multivariate issue that we dealt with is the multivariate, nonlinear dependence of the abundance of PM<sub>2.5</sub> in the atmospheric boundary layer on AOD, humidity, temperature, boundary-layer height, surface pressure, wind speed, and surface type. As mentioned earlier, our previous work in this area has won wide recognition as ground breaking.

## Method

NASA has a constellation of satellites flying in close formation called the “A-Train” (Figure 1). Several of these satellites host instruments that make a variety of aerosol observations. These instruments include Terra MODIS (Remer *et al.* 2005) and MISR (Kahn *et al.* 2005), launched in December, 1999; Aqua MODIS, launched in May, 2002; Aura OMI (Torres *et al.* 2007), launched in July, 2004; and CALIPSO CALIOP (Mcgill *et al.* 2007; Winker *et al.* 2007), launched in April, 2006. We also have aerosol observations from SeaWIFS (Hooker and McClain 2000), launched in August, 1997, on GeoEye's OrbView-2 satellite.

The aerosol optical depth (AOD),  $\tau$ , is a measure of the light extinction at a given wavelength by atmospheric aerosols, in a vertical column from the earth's surface up to the top of the atmosphere. Several of the A-Train instruments provide a daily global picture of the total aerosol optical depth. For example, MODIS provides the total AOD across its swath at a resolution of 10 km; the SeaWIFS resolution is 1.1 km. A new MODIS product at 3 km resolution should soon be available. The 3 km product introduces more noise but does capture fine (more urban scale) aerosol structure that is missed by the 10 km product. MODIS, OMI, and SeaWIFS provide the total global aerosol burden but not how it is distributed vertically, whereas other instruments provide detailed vertical aerosol structure but do not provide the contiguous global coverage of MODIS, OMI, and SeaWIFS. For instance, while CALIPSO provides corrected backscatter and extinction profiles at a 120 m vertical resolution, at altitudes below 20 km it does not provide contiguous horizontal coverage. MISR also provides some vertical information for cases with higher optical depths and distinct plume boundaries but at a coarser resolution than CALIPSO. The CALIPSO observations provide a set of high vertical resolution “curtains” underneath the satellite flight path. The CALIPSO curtains span the globe daily; however, there are substantial gaps between these curtains. Since CALIPSO completes 14.55 orbits per day, at the equator there is a separation of 24.7° in longitude between each successive curtain.

## Relating Aerosol Extinction to PM<sub>2.5</sub> Abundance

The relationship between the PM<sub>2.5</sub> abundance at the earth's surface and the boundary layer optical depth or aerosol extinction depends on a variety of factors that change both seasonally and geographically. These factors include the humidity, temperature, boundary-layer height, surface pressure, wind speed, and surface type (Liu *et al.* 2004a; Liu *et al.* 2004b; Hutchison *et al.* 2005; Gupta *et al.* 2006; Koelemeijer *et al.* 2006; Liu *et al.* 2007a; Liu *et al.* 2007b; Liu *et al.* 2007c; Pelletier *et al.* 2007; Gupta and Christopher 2008; Hutchison *et al.* 2008; Zhang *et al.* 2009).

When using a multi-linear analysis of the relationship between the AOD observed by MODIS and PM<sub>2.5</sub> it is found that better correlations are observed principally over the eastern United States in summer and fall (Zhang *et al.* 2009). The southeastern


United States has the highest correlation coefficients, at more than 0.6. The southwestern United States has the lowest correlation coefficient, at approximately 0.2. Several factors are at work here. One is that the entire aerosol loading does not usually reside in the boundary layer; hence, using AOD alone as a proxy for PM<sub>2.5</sub> will invariably result in significant error. For example, on the West Coast, a significant fraction of the AOD is due to smoke events where substantial amounts of aerosol are above the boundary layer. Additional reasons for the poor correlation in the southwest may be associated with the humidity and land surface type. In addition, the correlation depends on the version of the satellite retrieval. For example, MODIS v5.2.6 AOD retrievals demonstrate better correlation with PM<sub>2.5</sub> than v4.0.1 retrievals, but they have much less coverage because of the differences in the cloud-screening algorithm (Zhang *et al.* 2009). We address these issues by using a fully non-linear, multivariate, non-parametric machine learning approach.

(Gupta *et al.* 2006) found that correlation between AOD and PM<sub>2.5</sub> increases as the mixing-layer height decreases. Larger wind speed can induce high mixing-layer height, which can change the correlation between AOD and PM<sub>2.5</sub>. The relative humidity (RH) can affect the AOD-PM<sub>2.5</sub> by altering the optical properties of the aerosols. The higher the relative humidity, the larger the portion of light that is scattered, hence the larger AOD (Hoff and Christopher 2009). We address this issue in this study by using the humidity and boundary-layer height contemporaneously with each observation used. The humidity comes from the meteorological analyses. The boundary-layer height also derives from the meteorological analyses and can be verified with the available LIDAR data. The meteorological analyses we use are the NASA Modern Era Retrospective Analysis for Research and Applications (MERRA) analyses produced by the Goddard Space Flight Center (GSFC) Global Modeling and Assimilation Office (GMAO).

The correlation between AOD and PM<sub>2.5</sub> is also related to the surface pressure and wind speed (Smirnov *et al.* 1995; Lyamani *et al.* 2006; Choi *et al.* 2008; Rajeev *et al.* 2008). We address this issue by using the surface pressure and wind speed contemporaneously with each observation. The surface pressure and wind speed also come from the meteorological analyses.

**Table 1. Training dataset statistics and global 2000-2012 correlation coefficients.**

	n	R	R <sup>2</sup>
<b>Aqua Deep Blue</b>	8,233	0.99	0.98
<b>Aqua Standard</b>	30,298	0.99	0.98
<b>Terra Deep Blue</b>	4,011	0.99	0.98
<b>Terra Standard</b>	19,718	0.98	0.97



**Figure 1** Validation scatter diagrams showing the performance of the machine-learning algorithm for the two MODIS sensors using the standard and Deep Blue algorithms. In each case the x-axis shows the observed abundance of PM<sub>2.5</sub> (µg/cm<sup>3</sup>) as observed by in-situ instruments. The y-axis shows the abundance of PM<sub>2.5</sub> (µg/cm<sup>3</sup>) estimated by the machine learning based on the satellite and meteorological data products.

Several studies have sought to overcome this limitation by using satellite-derived Aerosol Optical Depth (AOD) with regression and/or numerical models to estimate ground-level PM<sub>2.5</sub> within the Earth's boundary layer. *Zhang et al.* (2009) presented a comprehensive study for the 10 EPA regions across the United States using multi-linear regression between the PM<sub>2.5</sub> abundance observed by the EPA and the Moderate Resolution Imaging Spectroradiometer (MODIS) AOD and a set of meteorological parameters. The best correlations of PM<sub>2.5</sub> with AOD were observed for the eastern states in summer and fall, with EPA region 4 having a correlation coefficient of more than 0.6. The poorest correlations were observed for the southwestern states, with EPA region 9 having a correlation coefficient of approximately 0.2. *Weber et al.* (2010) extended the study of *Zhang et al.* (2009) for five EPA monitoring sites in the Baltimore/Washington DC Metro area by considering AOD from MODIS, the Multi-Angle Imaging Spectroradiometer (MISR),



and the Geostationary Operational Environmental Satellite (GOES). The PM<sub>2.5</sub> estimates of *Zhang et al.* (2011) and *Weber et al.* (2010) are made available through the Infusing satellite Data into Environmental Applications (IDEA) website (<http://www.star.nesdis.noaa.gov/smcd/spb/aq/>).


In an elegant study *Van Donkelaar et al.* (2006) presented a global estimate of the long-term average PM<sub>2.5</sub> concentrations between 2001-2006 using both satellite observations of AOD from MODIS and a global chemical transport model to estimate  $\eta = \text{PM}_{2.5} / \text{AOD}$ . The 3D chemical transport model used was GEOS-Chem. *Van Donkelaar et al.* (2006) found significant spatial agreement with North American PM<sub>2.5</sub> measurements (correlation coefficient of 0.77) and with non-coincident measurements elsewhere (correlation coefficient of 0.83).

In this study we have used a proprietary machine learning approach to estimate  $\eta = \text{PM}_{2.5} / \text{AOD}$  entirely from observations. We used PM<sub>2.5</sub> observations from the United States, Europe, Africa, Australia and Asia to create a comprehensive training dataset spanning more than a decade. We then used this training dataset to estimate  $\eta$  as a function of the satellite AOD at multiple wavelengths and all the associated parameters that are available with the AOD (such as the angstrom exponent, scattering angle, cloud masks, surface reflectivity, and viewing geometry) and the meteorological analyses. Fifty independent trainings were performed using this training dataset, for each of these fifty trainings there was a random selection of 66% of the data for use in the training, with 34% of the data left out. The statistics shown in Table 1 and Figure 1 is the mean solution for these fifty independent trainings. Very careful attention is paid to ensure that the PM<sub>2.5</sub> observations and satellite observations are coincident in space and time to within a great circle separation of 0.02° (approximately 2 km) and a time window of 30 minutes. This is done for the standard and Deep Blue retrieval algorithms of MODIS Terra and Aqua. This can be thought of as the global fully non-linear multivariate extension to the pioneering work of (*Zhang et al.*, 2009).

The results of this comprehensive training are shown in the table below. The performance of the approach we have used here is substantially better than that of the previous studies. Our worst performance has a correlation coefficient of 0.85, which is better than the best performance of the previous studies 0.83 for the non-coincident measurements of *Van Donkelaar et al.* (2006). It should also be noted that our values are global, so include the west coast of the United States which, as mentioned above, is typically more challenging to reproduce (*Zhang et al.*, 2009).

As can be seen from Table 1 and Figure 1, we successfully used machine learning to describe the multivariate relationship between PM<sub>2.5</sub> and a suite of parameters including AOD. Example PM<sub>2.5</sub> distributions are shown in Figure 2. These daily distributions can be used to provide the time evolution of PM<sub>2.5</sub> exposure for individual personnel (e.g. Figure 3).

## PM<sub>2.5</sub> Monthly Average 2013/01



## PM<sub>2.5</sub> Monthly Average 2013/01




Figure 2. Example distributions of Monthly Average PM<sub>2.5</sub> ( $\mu\text{g}/\text{m}^3$ ) for January 2013.

## Los Angeles-Long Beach-Santa Ana, United States Population 0.8 Million




Figure 3. The Time evolution of PM<sub>2.5</sub> can be provided for more than a decade. The example above is for Long Beach, Los Angeles, CA.

### Key Research Accomplishments and Reportable Outcome

The key accomplishment of this study has been successfully using machine learning to provide daily global analyses of PM<sub>2.5</sub> from March 2000 up until the present. This is twice the length of the period we promised in the proposal. The fidelity of this analysis (as can be seen from Figure 1 and table 1) is significantly better than that of previous studies (*Van Donkelaar et al., 2006, Zhang et al., 2009, Zhang et al., 2011*). These PM<sub>2.5</sub> analyses are of considerable value to Global Health Surveillance (GHS), providing a capability to routinely estimate troop deployment exposure to elevated levels of particulate matter (PM) globally, significantly contributing to DoD-wide force health protection initiatives.

## References

- Anderson, T. L., R. J. Charlson, et al. (2003). "Mesoscale variations of tropospheric aerosols." Journal of the Atmospheric Sciences **60**(1): 119-136.
- Brown, M., D. Lary, et al. (2006). "Using neural nets to derive sensor-independent climate quality vegetation data based on AVHRR, spot-vegetation, seawifs and MODIS." Eos Trans. AGU **87**(36).
- Brown, M. E., D. J. Lary, et al. (2008). "Neural networks as a tool for constructing continuous NDVI time series from AVHRR and MODIS." International Journal of Remote Sensing **29**(24): 7141-7158.
- Campbell, J. R., J. S. Reid, et al. (2009). "CALIOP aerosol subset processing for global aerosol transport model data assimilation." J of Sel. Topics in Appl. Earth Obs. and Rem. Sens. **in press**.
- Charles, L., B. Gross, et al. (2007). Atmospheric transport of smoke and dust particulates and their interaction with the planetary boundary layer as observed by multiwavelength lidar and supporting instrumentation. Lidar remote sensing for environmental monitoring viii. U. N. Singh. **6681**: U157-U167.
- Choi, Y. S., C. H. Ho, et al. (2008). "Spectral analysis of weekly variation in pm10 mass concentration and meteorological conditions over china." Atmospheric Environment **42**(4): 655-666.
- Christensen, J. H. (1997). "The Danish eulerian hemispheric model - a three-dimensional air pollution model used for the arctic." Atmospheric Environment **31**(24): 4169-4191.
- Colarco, P. R., M. R. Schoeberl, et al. (2004). "Transport of smoke from canadian forest fires to the surface near washington, d. C.: Injection height, entrainment, and optical properties." Journal of Geophysical Research-Atmospheres **109**(D6).
- Colarco, P. R., O. B. Toon, et al. (2003a). "Saharan dust transport to the caribbean during pride: 1. Influence of dust sources and removal mechanisms on the timing and magnitude of downwind aerosol optical depth events from simulations of in situ and remote sensing observations." Journal of Geophysical Research-Atmospheres **108**(D19).
- Colarco, P. R., O. B. Toon, et al. (2003b). "Saharan dust transport to the caribbean during pride: 2. Transport, vertical profiles, and deposition in simulations of in situ and remote sensing observations." Journal of Geophysical Research-Atmospheres **108**(D19).
- Delfino, R. J., S. Brummel, et al. (2009). "The relationship of respiratory and cardiovascular hospital admissions to the southern california wildfires of 2003." Occupational and Environmental Medicine **66**(3): 189-197.
- Escudero, M., A. Stein, et al. (2006). "Determination of the contribution of northern africa dust source areas to pm10 concentrations over the central iberian peninsula using the hybrid single-particle lagrangian integrated trajectory

- model (hysplit) model." Journal of Geophysical Research-Atmospheres **111**(D6).
- Feldman, D. R., T. S. L'ecuyer, et al. (2008). "Remote sensing of tropical tropopause layer radiation balance using a-train measurements." Journal of Geophysical Research-Atmospheres **113**(D21).
- Gupta, P. and S. A. Christopher (2008). "Seven year particulate matter air quality assessment from surface and satellite measurements." Atmospheric Chemistry and Physics **8**(12): 3311-3324.
- Gupta, P., S. A. Christopher, et al. (2006). "Satellite remote sensing of particulate matter and air quality assessment over global cities." Atmospheric Environment **40**(30): 5880-5892.
- Hoff, R. M. and S. A. Christopher (2009). "Remote sensing of particulate pollution from space: Have we reached the promised land?" Journal of the Air & Waste Management Association **59**(6): 645-675.
- Hoff, R. M., J. A. Engel-Cox, et al. (2006). 3D-aqs, a three-dimensional air quality system - art. No. 629901. Remote sensing of aerosol and chemical gases, model simulation / assimilation, and applications to air quality. A. Chu, J. Szykman and S. Kondragunta. **6299**: 29901-29901.
- Hoff, R. M., S. P. Palm, et al. (2005). "Glas long-range transport observation of the 2003 california forest fire plumes to the northeastern us." Geophysical Research Letters **32**(22).
- Hogan, T. F. and T. E. Rosmond (1991). "The description of the navy operational global atmospheric prediction systems spectral forecast model." Monthly Weather Review **119**(8): 1786-1815.
- Hooker, S. B. and C. R. McClain (2000). "The calibration and validation of seawifs data." Progress in Oceanography **45**(3-4): 427-465.
- Houtekamer, P. L. and H. L. Mitchell (1998). "Data assimilation using an ensemble kalman filter technique." Monthly Weather Review **126**(3): 796-811.
- Hsu, N. C., S. C. Tsay, et al. (2006). "Deep blue retrievals of asian aerosol properties during ace-asia." Ieee Transactions on Geoscience and Remote Sensing **44**(11): 3180-3195.
- Huang, J., P. Minnis, et al. (2008). "Long-range transport and vertical structure of asian dust from calipso and surface measurements during pacdex." Journal of Geophysical Research-Atmospheres **113**(D23).
- Huang, J. P., P. Minnis, et al. (2007). "Summer dust aerosols detected from calipso over the tibetan plateau." Geophysical Research Letters **34**(18).
- Hutchison, K. D., S. J. Faruqui, et al. (2008). "Improving correlations between MODIS aerosol optical thickness and ground-based pm2.5 observations through 3D spatial analyses." Atmospheric Environment **42**(3): 530-543.
- Hutchison, K. D., S. Smith, et al. (2005). "Correlating MODIS aerosol optical thickness data with ground-based pm2.5 observations across texas for use in a real-time air quality prediction system." Atmospheric Environment **39**(37): 7190-7203.
- Jacquemin, B., J. Sunyer, et al. (2009). "Association between modelled traffic-related air pollution and asthma score in the ecrhs." European Respiratory Journal **34**(4): 834-842.

- Kahn, R. A., B. J. Gaitley, et al. (2005). "Multiangle imaging spectroradiometer (misr) global aerosol optical depth validation based on 2 years of coincident aerosol robotic network (aeronet) observations." Journal of Geophysical Research-Atmospheres **110**(D10).
- Karyampudi, V. M., S. P. Palm, et al. (1999). "Validation of the saharan dust plume conceptual model using lidar, meteosat, and ecmwf data." Bulletin of the American Meteorological Society **80**(6): 1045-1075.
- Koelemeijer, R. B. A., C. D. Homan, et al. (2006). "Comparison of spatial and temporal variations of aerosol optical thickness and particulate matter over europe." Atmospheric Environment **40**(27): 5304-5315.
- Kovacs, T. (2006). "Comparing MODIS and aeronet aerosol optical depth at varying separation distances to assess ground-based validation strategies for spaceborne lidar." Journal of Geophysical Research-Atmospheres **111**(D24).
- Lary, D. (2004). "Representativeness uncertainty in chemical data assimilation highlight mixing barriers." Atmospheric Science Letters **5**(1): 35-41.
- Lary, D. J. (2009). Artificial intelligence in geoscience and remote sensing. Aerospace technologies advancements.
- Lary, D. J. and O. Aulov (2008). "Space-based measurements of HCl: Intercomparison and historical context." Journal of Geophysical Research-Atmospheres **113**(D15).
- Lary, D. J. and L. Lait (2006). "Using probability distribution functions for satellite validation." Ieee Transactions on Geoscience and Remote Sensing **44**(5): 1359-1366.
- Lary, D. J., M. D. Muller, et al. (2004). "Using neural networks to describe tracer correlations." Atmospheric Chemistry and Physics **4**: 143-146.
- Lary, D. J., L. A. Remer, et al. (2009). "Machine learning and bias correction of MODIS aerosol optical depth." IEEE Geoscience and Remote Sensing Letters **6**(4): 694-698.
- Lary, D. J., D. W. Waugh, et al. (2007). "Variations in stratospheric inorganic chlorine between 1991 and 2006." Geophysical Research Letters **34**(21).
- Leptoukh, G., V. Zubko, et al. (2007). Spatial aspects of multi-sensor data fusion: Aerosol optical thickness. Igarss: 2007 ieee international geoscience and remote sensing symposium, vols 1-12 - sensing and understanding our planet: 3119-3122.
- Levy, R. C., G. G. Leptoukh, et al. (2009). "A critical look at deriving monthly aerosol optical depth from satellite data." Ieee Transactions on Geoscience and Remote Sensing **47**(8): 2942-2956.
- Liu, Y., M. Franklin, et al. (2007a). "Using aerosol optical thickness to predict ground-level pm2.5 concentrations in the st. Louis area: A comparison between misr and MODIS." Remote Sensing of Environment **107**(1-2): 33-44.
- Liu, Y., P. Koutrakis, et al. (2007b). "Estimating fine particulate matter component concentrations and size distributions using satellite-retrieved fractional aerosol optical depth: Part 1 - method development." Journal of the Air & Waste Management Association **57**(11): 1351-1359.
- Liu, Y., P. Koutrakis, et al. (2007c). "Estimating fine particulate matter component concentrations and size distributions using satellite-retrieved fractional

- aerosol optical depth: Part 2 - a case study." Journal of the Air & Waste Management Association **57**(11): 1360-1369.
- Liu, Y., R. J. Park, et al. (2004a). "Mapping annual mean ground-level pm2.5 concentrations using multiangle imaging spectroradiometer aerosol optical thickness over the contiguous united states." Journal of Geophysical Research-Atmospheres **109**(D22): 10.
- Liu, Y., J. A. Sarnat, et al. (2004b). "Validation of multiangle imaging spectroradiometer (misr) aerosol optical thickness measurements using aerosol robotic network (aeronet) observations over the contiguous united states." Journal of Geophysical Research-Atmospheres **109**(D6): 9.
- Liu, Z., D. Liu, et al. (2008). "Airborne dust distributions over the tibetan plateau and surrounding areas derived from the first year of calipso lidar observations." Atmospheric Chemistry and Physics **8**(16): 5045-5060.
- Lyamani, H., F. J. Olmo, et al. (2006). "Atmospheric aerosols during the 2003 heat wave in southeastern spain i: Spectral optical depth." Atmospheric Environment **40**(33): 6453-6464.
- Matichuk, R. I., P. R. Colarco, et al. (2007). "Modeling the transport and optical properties of smoke aerosols from african savanna fires during the southern african regional science initiative campaign (safari 2000)." Journal of Geophysical Research-Atmospheres **112**(D8).
- Matichuk, R. I., P. R. Colarco, et al. (2008). "Modeling the transport and optical properties of smoke plumes from south american biomass burning." Journal of Geophysical Research-Atmospheres **113**(D7).
- Mcgill, M. J., M. A. Vaughan, et al. (2007). "Airborne validation of spatial properties measured by the calipso lidar." Journal of Geophysical Research-Atmospheres **112**(D20).
- Pelletier, B., R. Santer, et al. (2007). "Retrieving of particulate matter from optical measurements: A semiparametric approach." Journal of Geophysical Research-Atmospheres **112**(D6): 18.
- Rajeev, K., K. Parameswaran, et al. (2008). "Observational evidence for the radiative impact of indonesian smoke in modulating the sea surface temperature of the equatorial indian ocean." Journal of Geophysical Research-Atmospheres **113**(D17).
- Reid, J. S., E. J. Hyer, et al. (2009). "Global monitoring and forecasting of biomass-burning smoke: Description of and lessons from the fire locating and modeling of burning emissions (flambe) program." Ieee Journal of Selected Topics in Applied Earth Observations and Remote Sensing **2**(3): 144-162.
- Remer, L. A., Y. J. Kaufman, et al. (2005). "The MODIS aerosol algorithm, products, and validation." Journal of the Atmospheric Sciences **62**(4): 947-973.
- Scholkopf, B., A. J. Smola, et al. (2000). "New support vector algorithms." Neural Computation **12**(5): 1207-1245.
- Sioutas, C., R. J. Delfino, et al. (2005). "Exposure assessment for atmospheric ultrafine particles (ufps) and implications in epidemiologic research." Environmental Health Perspectives **113**(8): 947-955.

- Smirnov, A., Y. Villevalde, et al. (1995). "Aerosol optical depth over the oceans - analysis in terms of synoptic air-mass types." Journal of Geophysical Research-Atmospheres **100**(D8): 16639-16650.
- Smola, A. J. and B. Scholkopf (2004). "A tutorial on support vector regression." Statistics and Computing **14**(3): 199-222.
- Torres, O., A. Tanskanen, et al. (2007). "Aerosols and surface UV products from ozone monitoring instrument observations: An overview." Journal of Geophysical Research-Atmospheres **112**(D24).
- Van Donkelaar, A., R. V. Martin, et al. (2006). "Estimating ground-level pm2.5 using aerosol optical depth determined from satellite remote sensing." Journal of Geophysical Research-Atmospheres **111**(D21).
- Vapnik, V. N. (1995). The nature of statistical learning theory. New York, Springer.
- Vapnik, V. N. (1998). Statistical learning theory. New York, Wiley.
- Winker, D. M., W. H. Hunt, et al. (2007). "Initial performance assessment of caliop." Geophysical Research Letters **34**(19).
- Xian, P., J. S. Reid, et al. (2009). "Impact of modeled versus satellite measured tropical precipitation on regional smoke optical thickness in an aerosol transport model." Geophysical Research Letters **36**.
- Yu, H., M. Chin, et al. (2010). "Global view of aerosol vertical distributions from calipso lidar measurements and gocart simulations: Regional and seasonal variations." Journal of Geophysical Research-Atmospheres.
- Weber, S. A., Engel-Cox, J. A., Hoff, R. M., Prados, A. I. & Zhang, H. (2010) An improved method for estimating surface fine particle concentrations using seasonally adjusted satellite aerosol optical depth. *Journal of the Air & Waste Management Association*, 60, 574-585.
- Zhang, H., R. M. Hoff, et al. (2009). "The relation between moderate resolution imaging spectroradiometer (MODIS) aerosol optical depth and pm2.5 over the united states: A geographical comparison by us environmental protection agency regions." Journal of the Air & Waste Management Association **59**(11): 1358-1369.
- Zhang, J. L., J. S. Reid, et al. (2008). "A system for operational aerosol optical depth data assimilation over global oceans." Journal of Geophysical Research-Atmospheres **113**(D10).
- Zhang, H., Lyapustin, A., Wang, Y., Kondragunta, S., Laszlo, I., Ciren, P. & Hoff, R. M. (2011) A multi-angle aerosol optical depth retrieval algorithm for geostationary satellite data over the united states. *Atmospheric Chemistry and Physics*, 11, 11977-11991.
- Zubko, V., G. Leptoukh, et al. (2009). "Study of data merging and interpolation methods for use in an interactive online analysis system: MODIS terra and aqua daily aerosol case." IEEE Transactions on Geoscience and Remote Sensing **submitted August 2009**.